

DAUWE PLASTIC SURGERY

GENERAL INSTRUCTIONS PLEASE READ THIS VERY CAREFULLY

PRE-OP VISIT

- At your pre-op visit you can expect:
 - ☐ Blood work ☐ EKG (>50 years old) ☐ Sign consents ☐ Receive Medications **Rx**
- Make sure all balances are paid. If your health insurance is helping pay for surgery, you will be responsible for your deductible and co-insurance.
- Make sure you have all your medications filled before surgery day.
- **Do not use any nicotine products 4 weeks before and 4 weeks after surgery.** This includes patches, gums, and e-cigarettes.
- Avoid alcoholic beverages for 1 week before and 1 week after surgery.
- Avoid any medications that thin your blood for 1 week before and 3 weeks after surgery (ibuprofen, naproxen, aspirin, etc.).
- If you are taking a prescription blood thinner (aspirin, plavix, Coumadin, xarelto, etc.), please let Dr. Dauwe know, because this will need to be stopped before surgery.
- If you have any serious health conditions, please discuss the need for preoperative health clearance with Dr. Dauwe before surgery.

THE MORNING OF SURGERY

- **Wake up 4 hours before surgery and drink 20oz of Gatorade. Do not eat or drink anything else. Nothing else to eat or drink 8 hours prior to surgery.** (If you have diabetes, this does not apply to you.)
- Do not put on make up, lotions, creams, or other cosmetic products.
- Remove all piercings and jewelry.
- For surgeries involving implants, on the morning before surgery wash your body from neck down with Hibiclens (chlorhexidine 2%) soap.

WHAT TO EXPECT AFTER SURGERY

- Stay active after surgery. This prevents blood clots from forming in your legs following surgery.
- Keep your **heart rate** below 100 for 3 weeks and below 130 for 6 weeks.
- Avoid **alcoholic beverages** for 1 week after surgery. It will make you bruise and perhaps bleed after surgery.
- Do not take any medications that **thin your blood** (ibuprofen, aspirin, naproxen) for 1 week before and 3 weeks after surgery.
- Take your **pain medication** every 4-6 hours for the first 24 hours, then as needed.
- Narcotics cause constipation, so please try to wean yourself from these meds as soon as possible. Eat high fiber foods and take a stool softener like Senna or Colace (over the counter).
- If you are constipated, please follow this regimen of medications:

Milk of Magnesia → Dulcolax Suppository → Magnesium Citrate

DAUWE PLASTIC SURGERY

- You may take Tylenol, but some pain medications have Tylenol (acetaminophen) in them. Do not take more than 4000mg of Tylenol in a 24-hour period.
- You may **drive** after you are off narcotic pain medication, and you are able to physically react to traffic (swerving or slamming on your brakes).
- Sensations like numbness, tingling, and burning are common during the healing process. These sensations may last several weeks and will gradually disappear.
- Bruising and swelling are normal for 2-3 weeks. Sometimes this lasts longer, but it always goes away.
- You may **shower**, but face away from the stream of water for first 7 days.
- Remove your steri-strips at 2 weeks.
- If you have been provided with compression garments, wear these for 6 weeks following surgery.
- Do not bath/swim/submerge after surgery until Dr. Dauwe clears you. This is a serious infection risk.

TO SPEED YOUR RECOVERY

- ✓ **Take a nap.** After you leave our facility, go home and take a 2-hour nap. This will help you recover from the anesthetic and get you back in your home routine.
- ✓ **Eat something substantial.** You haven't eaten since the day before surgery. Your empty stomach is contributing to your fatigue and discomfort. Eat a normal meal, and you will feel better.
- ✓ **Stay active, but not too active.** Stay out of bed most of the day and move around the house. Keep your heart rate below 100 though.
- ✓ **Minimize narcotic pain medication.** Narcotic pain medication sedates you and slows you down. It will make you want to lie down. When you lie down, you get bored, and you think about how much you are hurting.
- ✓ **Distract your brain.** If you engage activities that distract you from the fact that you just had surgery, you will recover faster. Do not exert yourself, but watch TV, sit at the kitchen table and eat lunch/dinner, speak with friends/family on the phone, and engage in conversation. This helps tremendously.
- ✓ **Move around.** Keep your joints loose and muscles stretched. The stiffness in your shoulders, elbows, neck, and back contributes to your discomfort. Just like walking on a sprained ankle, moving around after surgery will quicken your recovery.

THINGS TO LOOK OUT FOR

If you have any of the following, please call the office:

- ✗ vomiting ✗ rash ✗ shortness of breath ✗ chest pain ✗ leg swelling ✗ diarrhea
- ✗ fever ($>100.5^{\circ}$) ✗ redness ✗ increasing tenderness at the surgical site

DAUWE PLASTIC SURGERY

IF YOU HAVE ANY QUESTIONS, PLEASE CALL THE OFFICE

- It is important to be seen according to the follow up timeline. Dr. Dauwe will explain your postoperative needs at each follow up visit.
- The office is open 9-5p (Mon-Thurs) and 9-3p (Fri).
- If you need immediate care, please call Dr. Dauwe's office at anytime. There is no such thing as a silly question. I would rather you call me and get it right, than not "bother" me and chance not doing the right thing.
- If you had any kind of implants placed and have a dental procedure or major surgery planned, please notify Dr. Dauwe, so he can prescribe a pre-operative antibiotic to protect your implants from possible infection.
- Please call the office and schedule the first follow up appointment.


SPECIFIC INSTRUCTIONS FOR EACH SURGERY

If you have drains

- Keep the drain compressed. The drain uses negative pressure to pull fluid out of the surgery site
- Remove the cap from the top of the drain, empty the drain completely, record the daily total, compress the drain, and replace the cap
- If you have implants, you must be on antibiotics while drains are in place

Breast Implants

- Do not submerge your incisions under water for 6 weeks
- If your implants were repositioned it is normal to have a "dent" or contour irregularity on the side of your breast. This is the internal stitch line and always smooths out
- You may remove your ACE wrap on day 2 and put on a soft, elastic bra (no underwire)
- Dr. Dauwe likes the brand "True & Co" for recovery bras (see picture)
- Please do not do any strong upper body exercises for 3 months (bench press, pushup, inverted yoga, etc.)
- If you had an operation for capsular contracture, you will lose the upper pole fullness/roundness from the contracted implant. Your new breast will be softer and more natural.



Breast Reduction or Lift

- You may remove your ACE wrap on day 2 and put on a soft, elastic bra (no underwire)
- Dr. Dauwe likes the brand "True & Co" for recovery bras
- You should wait 3 months to buy new bras, as your breasts will change over that time

Tummy Tuck

- **VERY IMPORTANT: YOU MUST WALK AFTER SURGERY.** This prevents blood clots from forming in your legs and a potentially fatal pulmonary embolus
- If you notice any swelling in your feet/legs or shortness of breath, call Dr. Dauwe immediately

DAUWE PLASTIC SURGERY

- It is normal to walk bent over for the first week following surgery due to muscle spasm. Valium (diazepam) will help with muscle spasm and discomfort
- Wear your compression garment (abdominal binder) for 6 weeks postop
- Any dimples in your tummy skin will go away. That is the internal stitches securing your skin down to the muscle

Liposuction

- Your liposuction incisions will leak blood-colored fluid. It is best to sleep on a few towels, so this fluid does not stain your mattress
- It is normal to feel firm areas and areas of unequal swelling. As the swelling goes away, the areas will even out
- It takes a full year for the swelling to go away completely

Face/Neck Lift

- It is normal to feel tight and swollen and mild discomfort. Severe pain is not normal, and you should discuss with Dr. Dauwe if you are experiencing this
- Numbness in your cheek, neck, and ears is normal and will improve with time
- Stitches will be removed at one week. Sometimes, we will leave a few stitches longer to minimize scarring

Eyelid Surgery

- Bruising and swelling will be more dramatic around your eyes because the eyelid skin is so thin. Occasionally, your eyes will swell shut
- Use your prescription drops as prescribed, and use artificial tears (over the counter) if your eyes are feeling dry
- If you notice your eye(s) bulging, severe pain, or visual changes, please call Dr. Dauwe
- Stitches will be removed at one week

DAUWE PLASTIC SURGERY

RECOVERY AND FOLLOW UP TIMELINE

One Day

You will be feeling tired and sore. The bruising and swelling are the worst 2-3 days following surgery. Take your pain meds as needed, take a few naps, but stay reasonably active. Avoid blood thinners, alcohol, and nicotine.

One Week

Your soreness will be improving, but your fatigue will persist. Dr. Dauwe will remove your stitches at this visit if necessary. Your bruising will be improving, but your swelling will persist. Try not to stay out of the house for longer than a few hours, because it will exhaust you.

Two - Three Weeks

Your soreness and fatigue have resolved. You may return to light to moderate exercise (HR>100). Your bruising has resolved and your swelling will be improving. Now you may drink alcoholic beverages. It is now ok to resume any blood thinners you may have been taking. Start scar therapy protocol.

Six Weeks

You will be feeling much better, and your results will really start to shine. You may return to intense exercise (HR>130), but if you had breast implant surgery, please do not do any strong upper body exercises yet (bench press, pushup, inverted yoga, etc.).

Three Months

Return to the office for postop photographs. At this point, 90% of the swelling has resolved and you are close to your final result. If you had breast implants, you may return to upper body strength exercises.

Six Months

At this visit, postop photographs will be repeated. You will see your final results now, as 95% of swelling has resolved.

One Year

Your postop photographs will be repeated, and you are finished with your surgical recovery. 100% of swelling has resolved.

Long-Term

You are a patient for life. If you have any concerns or interests, we will always be a resource for you.